

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12Q 1/68, C12P 19/34</b>	<b>A1</b>	(11) International Publication Number: <b>WO 00/01849</b> (43) International Publication Date: <b>13 January 2000 (13.01.00)</b>
<p>(21) International Application Number: <b>PCT/US99/15030</b></p> <p>(22) International Filing Date: <b>1 July 1999 (01.07.99)</b></p> <p>(30) Priority Data: <b>60/091,578</b>      <b>2 July 1993 (02.07.93)</b>      <b>US</b></p> <p>(71) Applicant (for all designated States except US): <b>INVITRO DIAGNOSTICS, INC. (US/US); Suite 407, 3960 Broadway, New York, NY 10032 (US).</b></p> <p>(72) Inventor; and (73) Inventor/Applicant (for US only): <b>GROSSMAN, Abraham (US/US); 666 Washington Avenue, Pleasantonville, NY 10570 (US).</b></p> <p>(74) Agent: <b>JANIUK, Anthony, J.; InVitro Diagnostics, Inc., Suite 407, 3960 Broadway, New York, NY 10032 (US).</b></p>	<p>(81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b></p> <p>Published With international search report.</p>	
<p>(54) Title: <b>METHODS, COMPOSITIONS AND APPARATUS FOR MAKING NUCLEIC ACID MOLECULES HAVING A SELECTED AFFINITY TO A TARGET MOLECULE</b></p>		
<p>(57) Abstract</p> <p>The present invention is directed to methods, compositions, kits and devices for making a nucleic acid having selected affinity to a target molecule. One embodiment of the present invention is a method of making a replicatable nucleic acid template having a selected affinity to a target. The method comprises the step of applying a selection to a first generation comprising at least one replicatable nucleic acid template as the replicatable nucleic acid template is replicated by a nucleic acid polymerase to form at least one subsequent generation comprising a replicatable nucleic acid template. The selection is based on the affinity of the replicatable nucleic template of different generations to the target. The nucleic acid polymerase introduces genetic variability between generations of the replicatable nucleic acid template to produce replicatable nucleic acid templates having different affinities to the target. The replicatable nucleic acid templates are separated on the basis of the affinity of the replicatable nucleic acid template to the target.</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RU	Romania		
CL	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Methods, Compositions and Apparatus for Making Nucleic Acid Molecules  
Having a Selected Affinity to a Target Molecule

5 Field of the Invention

The present invention is directed to methods, compositions, kits and devices for making a nucleic acid molecule having selected affinity to a target molecule. One embodiment of the present invention is a method of making a replicatable nucleic acid template having a selected affinity to a target. The method comprises the step of applying  
10 a selection to a first generation comprising at least one replicatable nucleic acid template as the replicatable nucleic acid template is replicated by a nucleic acid polymerase to form at least one subsequent generation. The subsequent generation comprises at least one evolved replicatable nucleic acid template. The selection is based on the affinity of the replicatable nucleic templates of different generations to the target. The nucleic acid  
15 polymerase introduces genetic variability between generations of the replicatable nucleic acid template, directed by selection, to produce evolved replicatable nucleic acid templates having different affinities to the target. The replicatable nucleic acid templates are separated on the basis of the affinity of the replicatable nucleic acid template to the target.

20 Background of the Invention

The words and terms used in this application have definitions and meanings that one skilled in the art of molecular biology and genetics would recognize. For the purpose of convenience, the following definitions are provided.

Nucleic acid means either deoxyribonucleic acid (DNA) or ribonucleic acid  
25 (RNA), single stranded or double stranded and any chemical modifications thereof.

A replicatable nucleic acid template is a nucleic acid having a sequence of nucleotides which sequence is recognized by a polymerase to produce copies of itself. By way of example, without limitation, certain RNA bacteriophages replicate a nucleic acid, an RNA, by an RNA-dependent RNA polymerase. The enzymes Q-Beta replicase and  
30 other replicases, such as MS2 replicase (Fedoroff, N. 1975 Cold Spring Harbor Symposium 235-258), GA replicase (Yousani et al 1981 J. Biochem 89,741-750), and SP replicase (Miyare et al 1971 PNAS 68, 2022-2024) are examples of such RNA polymerases. These polymerases typically exhibit preferences for sequences having certain features, particularly their respective naturally occurring sequences or templates.

The term "generation" refers to a group of replicatable nucleic acid templates initiating or resulting from a cycle of replication. The homogeneity of the population of

5 replicatable nucleic acid templates will depend on the fidelity of the polymerase in replicating the template. The enzyme Q-Beta replicase replicates RNA templates with low fidelity. Rather, each generation of templates will exhibit differences caused by a single nucleotide substitution, addition and deletion and the rearrangement of groups of nucleotides.

10 A ligand is a molecule which exhibits affinity for another molecule. A molecule of interest or target is that which one wishes to detect, identify and characterize in greater detail or create a nucleic acid ligand for. Affinity refers to the ability to bind to or form a complex with.

A kit is an assembly of parts and reagents for performing a method.

15 Other words and terms will be defined in the Summary of Invention and Detailed Discussion which follow.

Nucleic acids can exhibit affinity for other nucleic acids (other than Watson-Crick hybridization), proteins, carbohydrates, peptides, metabolites, and other small molecules. However, application of such nucleic acid molecules, as ligands, in the manner of  
20 antibodies, has been limited. Present techniques, for identifying nucleic acid ligands which exhibit an affinity for a chosen molecule of interest or target, are awkward. Present techniques require the synthesis of random libraries of nucleic acids. Present techniques require multiple reiterative steps.

For example, US Patent 5,270,163 requires the steps of contacting a library of  
25 randomized nucleic acid molecules with the target, partitioning the high affinity nucleic acid molecules from the low affinity nucleic acid molecules and amplifying the high affinity nucleic acid molecules. These steps are repeated as required. In this process, the nucleic acid molecules contacting the target is an RNA and the amplified nucleic acid is a cDNA of such RNA. The process is time consuming. Finally, one or more species of  
30 nucleic acid are identified with high affinity to the target from the many with low affinity to the target.

Embodiments of the present invention address several problems associated with identifying nucleic acid ligands. Rather than identifying one or a small number of species of nucleic acid exhibiting the highest affinity for the target, embodiments of the present

invention feature the evolution of replicative nucleic acid templates which over generations exhibit a selected affinity to the target.

5    **Summary of the Invention**

The present invention is directed to methods, compositions, kits and devices for making a nucleic acid molecule having selected affinity to a target molecule. One embodiment of the present invention is a method of making a replicatable nucleic acid template having a selected affinity to a target. The method comprises the step of applying  
10    a selection to a first generation comprising at least one replicatable nucleic acid template as the replicatable nucleic acid template is replicated by a nucleic acid polymerase to form at least one subsequent generation comprising a replicatable nucleic acid template. The selection is based on the affinity of the replicatable nucleic template of different generations to the target. The nucleic acid polymerase introduces genetic variability  
15    between generations of the replicatable nucleic acid template to produce replicatable nucleic acid templates having different affinities to the target. The replicatable nucleic acid templates are separated on the basis of the affinity of the replicatable nucleic acid template to the target.

As used herein, the term "selection" refers to conditions in which variants of the  
20    replicatable nucleic acid template compete. Each variant may present a unique molecular genotype which, due to its nucleic acid composition, will represent a unique phenotype. The conditions of the selection allow variants with a particular phenotype to replicate at a higher rate than others. The nucleic acid polymerase introduces molecular variability with each generation, to create or evolve phenotypes and genotypes not present in the initial  
25    generation.

A preferred nucleic acid is an RNA. And, a preferred RNA replicatable template is selected from the group of templates recognized by the nucleic acid polymerase. By way of example, without limitation, the enzyme Q-Beta replicase recognizes the following templates; wild-type Q-Beta phage genomic RNA, midi-varient RNA, nano-varient RNA,  
30    RQ-135 RNA and modifications of such RNAs derived from evolutionary processes.

A preferred polymerase is selected from the group of polymerases consisting of Q-Beta replicase, MS2 replicase, GA replicase, and SP replicase and modifications thereof. A particularly preferred replicase is a modified RNA polymerase exhibiting relaxed selectivity towards the template and relaxed fidelity in the replication process to create

variants. Such polymerase tends to introduce and accept nucleotide substitution, deletion and insertion at a high rate.

One embodiment of the present invention features an RNA polymerase having one or more target moieties. The RNA polymerase having one or more target moieties exhibits affinity to the target to exert a selection on the RNA template as such template is replicated. A preferred RNA polymerase carries one or more target moieties covalently bound to a unit or subunit of the enzyme. By way of example, without limitation, the enzyme Q-Beta replicase has four subunits; comprising a ribosomal protein S1, EF-Ta, EF-Tu, and a viral subunit. The target is bound to or incorporated in one or more subunits. Where the target is a protein the nucleic acid encoding such target is cloned into the nucleic acid encoding such subunit and expressed as part of such subunit.

For purposes of convenience, a non-naturally occurring RNA polymerase with one or more target moieties is referred to in this application as an "evolase". The evolase may have naturally occurring units and subunits to which the target is not normally bound. Replicable RNA templates with high affinity to the target tend to be replicated with greater frequency than replicable RNA templates with low affinity by an evolase in the presence of suitable replication conditions. In the context of the present application, the terms "high" and "low", as applied to affinity, refer to relative affinities. This preferential replication applies a selection for high affinity templates. The replicable RNA templates evolve to increase the relative number of high affinity templates over successive generations.

One embodiment of the present invention features selection applied by passing a solution comprising the first and subsequent generations of the replicable nucleic acid template through or over a fixed medium. The fixed medium has the target immobilized, such that replicable nucleic acid templates having low affinity for the target and replicable nucleic acid templates having high affinity for the target assume different positions in the solution as the solution passes through the fixed medium. The medium may comprise a capillary, packed column, a gel, beads, glass or plastic sheets and other particles placed in any suitable vessel.

Preferably, the fixed medium is provided with reagents and a polymerase. The reagents and polymerase are added to the fixed medium in a manner in which the replicable nucleic acid templates may be replicated to form at least one subsequent generation. Replicable nucleic acid templates with low affinity will assume a different

position in the solution. Such low affinity replicatable nucleic acid templates will pass more quickly, be withheld less, than such replicatable nucleic acid template with high affinity. Preferably, the fixed medium allows the formation of a multitude of generations in the presence of the target. Thus, replicatable nucleic acid templates exhibiting high or low affinity, or any affinity in between, can be created. The target can be fixed by itself or attached covalently to the nucleic acid polymerase. Both the nucleic acid polymerase and the target can be immobilized.

Preferably, the solution flows through a fixed matrix. Preferably, the solution and RNA templates are influenced or acted upon by electromotive forces. These electromotive forces, in the nature of electric field electrophoresis, move the templates through the matrix.

In the alternative, selection is applied by placing the first generation and target in a solution and forming at least one subsequent generation. The method further comprises the step of forming a first fraction of the solution and a second fraction of said solution. The first fraction is substantially devoid of target. The solution of said first fraction containing replicatable nucleic acid templates having lower affinity than replicatable nucleic acid templates of the second fraction. The second fraction has the target or was closely associated with the target. This fraction contains replicatable nucleic acid templates having higher affinity than replicatable nucleic acid templates in the first fraction. Preferably, the formation of fractions is repeated or is performed in a continuous manner during the formation of multiple generations. That is, at least one fraction containing a replicatable nucleic acid template of a subsequent generation is again placed with the target in a solution. After the formation of a further subsequent generation of a replicatable nucleic acid template, a further first and second fraction is formed. Thus, replicatable nucleic acid templates exhibiting high or low affinity, or any affinity in between, can be created.

Fractions may be formed by one or more of the processes selected from the group consisting of centrifugation, dilution, filtration, electric field electrophoresis, precipitation and immobilization of the target on supports. The supports are preferably removable, such as magnetic beads, styrene beads and particles, nitrocellulose and the like.

Preferably, the selection is applied in a plurality of vessels and/or in a plurality of media to reduce selection based on the composition of the vessel.



A further embodiment of the present invention features a device for making a replicatable nucleic acid template having a selected affinity to a target. The device comprises means for applying a selection to a first generation comprising at least one replicatable nucleic acid template as the replicatable nucleic acid template is replicated by a polymerase to form at least one subsequent generation comprising a replicatable nucleic acid template. The selection is based on the affinity of the replicatable nucleic acid template of different generations to the target. The polymerase introduces molecular variability between generations of the replicatable nucleic acid template to produce replicatable nucleic acid templates having different affinities to the target. And, the device comprises means for separating the replicatable nucleic acid templates on the basis of the affinity of the replicatable nucleic acid template to the target.

Preferably, the means for applying a selection is a fixed medium. The selection is applied by passing a solution comprising the first and subsequent generations of the replicatable nucleic acid template through the fixed medium. The fixed medium has the target immobilized and capable of being bound to the replicatable nucleic acid templates. Replicatable nucleic acid templates having low affinity for the target and replicatable nucleic acid templates having high affinity for the target assume different positions in the solution as the solution passes through the fixed medium.

Preferably, the fixed medium is held in a column.

In the alternative, the fixed medium is a matrix upon which an electromotive force is placed. Or, the means for applying a selection is a non-naturally occurring polymerase having a target moiety, a evolvasse.

In a further embodiment of the present invention, the means for applying a selective vector comprises a plurality of vessels constructed and arranged to receive fractions of solutions. The fractions contain replicatable nucleic acid templates exhibiting different affinities to the target. The replicatable nucleic acid templates exhibiting a trend toward the desired affinity are allowed to form at least one further generation.

A further embodiment of the present invention features a kit for making a replicatable nucleic acid template having a selected affinity to a target. The kit comprises means for applying a selection to a first generation comprising at least one replicatable nucleic acid template as the replicatable nucleic acid template is replicated by a polymerase to form at least one subsequent generation comprising a replicatable nucleic acid template. The selection is based on the affinity of the replicatable nucleic acid template of different

generations to the target. The polymerase introduces genetic variability between generations of the replicatable nucleic acid template to produce replicatable nucleic acid templates having different affinities to the target. And, the kit comprises means for  
5 separating the replicatable nucleic acid templates on the basis of the affinity of the replicatable nucleic acid template to the target.

A further embodiment of the present invention comprises an evolvase.

These and other features and advantages will be apparent to individuals skilled in the art in light of the drawings and the detailed description of the invention which follows.

10

#### Brief Description of the Drawings

Figure 1 depicts the nucleic acid sequence of one replicatable nucleic acid template for the enzyme Q-Beta replicase;

15 

Figure 2 depicts the enzyme having features of the present invention;

Figure 3 depicts in schematic form a device embodying features of the present invention;

Figure 4 depicts in schematic form a further device embodying features of the present invention;

20 

Figure 5 depicts in schematic form a further device embodying features of the present invention; and,

Figure 6 depicts a kit having features of the present invention.

#### Detailed Description of the Drawings

25 The present invention is directed to methods, kits and devices for making a nucleic acid having selected affinity to a target molecule. The invention and the figures will be described in detail with respect to a method of making a replicatable nucleic acid template having a selected affinity to a target. The method comprises the step of applying a selection to a first generation comprising at least one replicatable nucleic acid template as  
30 the replicatable nucleic acid template is replicated by a nucleic acid polymerase to form at least one subsequent generation comprising a replicatable nucleic acid template. The selection is based on the affinity of the replicatable nucleic template of different generations to the target. The nucleic acid polymerase introduces genetic variability between generations of the replicatable nucleic acid template to produce replicatable  
35 nucleic acid templates having different affinities to the target. The replicatable nucleic acid

templates are separated on the basis of the affinity of the replicatable nucleic acid template to the target.

A preferred nucleic acid is an RNA. One replicatable RNA template is midi-variant RNA, the sequence of which is set forth below in Seq ID No 1:

Seq ID No. 1

5'-GGGGACCCCC CCGGAAGGGG GGGACGAGGU GCGGGCACCU'  
CGUACGGGAG UUCGACCGUG ACGCUCUAGA GAUCUAGAGC  
ACGGGCUAGC GCUUUCGCGC UCUCUCCAGGU GACGCCUCGU  
GAAGAGGCGC GACCUUCGUG CGUUCGGUG ACGCACGAGA  
ACCGCCACGC UGCUUCGCAG CGUGGCUCUCCU UCGCGCAGCC  
CGCUGCGCGA GGUGACCCCC GAAGGGGGGU UCCC-3'

Figure 1 depicts in schematic form naturally occurring MDV(+) RNA. Nishiharra et al (1983). Other replicatable RNA templates may also be used depending on the choice of polymerase. With respect to the enzyme Q-Beta replicase, wild type replicatable RNA, Q-Betabeta cteriphage genomic RNA, may be preferred.

A preferred polymerase is selected from the group of polymerases consisting of Q-Beta replicase, MS2, GA, and SP and modifications thereof. A particularly preferred replicase is a modified RNA polymerase exhibiting relaxed selectivity towards the template and relaxed fidelity in the replication process to create molecular variants. A relaxed fidelity RNA replicase is described by Yoshio Inokuchi and Mayayuki Kajitani, "Deletion Analysis of QB Replicase," *J. Bio Chem.*, vol. 272, no. 24, 15339-15345 (1997). The variants of the replicatable nucleic acid template compete for replication by the polymerase, with each other and each prior generation of replicable nucleic acid templates.

One preferred enzyme, referred to herein as an evolvase, is represented by the formula below:



wherein C represents a catalytic entity and T represents a target. The target is covalently bound to the catalytic entity or bound by affinity. Methods and compositions for affinity coupling, such as strep-avidin, or immunological reactions are well-known. Where the target is a protein, the chemistry to covalently bind a protein to the catalytic entity is well-known. A preferred catalytic entity is selected from the group of polymerases consisting of Q-Beta replicase, MS2, GA, and SP and modifications thereof. A particularly preferred replicase is a modified RNA polymerase exhibiting relaxed fidelity in the replication process to create molecular variants. Figure 2 depicts an evolvase derived from Q-Beta replicase. As depicted, the evolvase comprises a ribosomal protein S1, EF-Ts, a viral subunit and in the place of a subunit, EF-Tu, a target protein subunit. A target subunit protein is formed by cloning sequences for the target into sequences for the subunit EF-Tu. In the alternative, the sequences encoding the target can be cloned into sequences for the other subunits. The cloned sequences are expressed and the evolvase isolated. In the further alternative, a polymerase is chemically altered to covalently attach the target to an existing enzyme through protein chemistry. The formula above is not intended to reflect a strict stoichiometric relationship between the catalytic entity and the target. Indeed, a plurality of target entities may be combined with each catalytic entity.

The evolvase is placed in contact with a nucleic acid template. The template is replicated by imposing reaction conditions. In the evolvase based on the enzyme Q-Beta replicase, the reaction conditions would be identical as those for such enzyme. The reagents comprise 173 mM Tris-HCl, pH 7.5, 27 mM MgCl<sub>2</sub>, and .77mM each ATP, GTP, CTP, and UTP. Upon imposing reaction conditions, successive generations of the template are formed. Molecular variability is introduced in the successive generations by the evolvase. Those templates with the greatest affinity to the target will form the dominant species. These dominant species may be isolated or serially diluted and reapplied to the evolvase for further evolution. The early generations will exhibit low affinity to the target. The latter generations will exhibit high affinity to the target.

In the alternative, a preferred selection is applied by passing a solution comprising the first and subsequent generations of the replicatable nucleic acid template through or over a fixed medium. The fixed medium has the target immobilized, such that replicatable  
5 nucleic acid templates having low affinity for the target and replicatable nucleic acid templates having high affinity for the target assume different positions in the solution as the solution passes through or over the fixed medium. The medium may comprise a capillary, packed column, a gel, beads, glass or plastic sheets, and other particles placed in any suitable vessel.

10 Turning now to Figure 3, a device for making a nucleic acid having selected affinity to a target molecule is illustrated in partial cut-away form. The device, generally designated by the numeral 11, comprises the following major elements, a housing 15 and a support 17. The housing 11 has a cylindrical shape having two ends 21 and 23. Housing 15 is preferably made of plastic or glass. As illustrated, one end 21 has a taper for retaining  
15 support 17. To further retain the support 17, the device 11 has a frit 25 retained in housing 15 at the tapered end 21. Tapered end 21 has an opening 31 to allow solutions to exit from the housing 15. The opposing end 23 is adapted to receive replicatable nucleic acid templates, a polymerase and reagents for replication.

The support 17, as illustrated, comprises a matrix such as Activated CH Sepharose 4B  
20 (Pharmacia LKB). The target, in the case of proteins, is attached to the matrix according to standard procedures. The activated groups remaining on the matrix are blocked with Tris.

In operation, the housing 15 is loaded with the support 17 to which the target is affixed. The support 17, a fixed medium, is provided with a replicatable nucleic acid  
25 template. A preferred template is Q-Beta bacteriophage genomic RNA. The template is preferably allowed to migrate into the support 17 matrix.

Reagents and a polymerase are added to the support 17 matrix. The reagents and polymerase are added to the support 17 in a manner in which the replicatable nucleic acid templates may be replicated to form at least one subsequent generation. Where the  
30 polymerase is Q-Beta replicase, the reagents comprise 173 mM Tris-HCl, pH 7.5, 27 mM  $MgCl_2$ , and .77 mM each ATP, GTP, CTP, and UTP. Where the housing 15 is sized to correspond approximately with the size of an Eppendorf tube, approximately 1.2  $\mu$ g Q-Beta replicase is added. As the reaction proceeds, the solution is drawn through the housing 15 through the support 17 by means of vacuum at end 21, pressure at end 23 or centripetal

force by means of a centrifuge (not shown). As the reaction proceeds, new generations of the template are formed. Due to the lack of template specificity and the lack of replication fidelity of the enzyme, the templates will evolve, losing or gaining or substituting single nucleotides or nucleic acid sequences, in response to the selection.

Replicable nucleic acid templates with low affinity will assume a different position in the solution. Such low affinity replicatable nucleic acid templates will pass more quickly, be withheld less, than such replicatable nucleic acid template with high affinity. Templates exhibiting high affinity will form more generations being held in the support for a longer period. Thus, replicatable nucleic acid templates exhibiting high or low affinity, or any affinity in between, can be created.

Templates with high affinity to the target can be subjected to further selection or modification to evolve such templates for low affinity to the support 17. That is, the device 11 is used with a support 17 which support 17 does not have target. Templates which pass through the support 17 first have low affinity for the support 17.

Figure 4 a and b depicts a further embodiment of the present invention. Turning first to Figure 4a, a device, generally designated by the numeral 41 for making a nucleic acid having selected affinity to a target molecule is illustrated. The device 41 comprises a first vessel 45. First vessel 45 receives a replicatable nucleic acid template, target, reagents and polymerase. Upon imposing reaction conditions, at least one subsequent generation of template is formed. The double arrow represents imposing fraction forming conditions on the target. These conditions may be imposed by chemical or physical means, such as the addition of salt, change in pH, of centrifugation. The imposition of fraction forming conditions forms a first fraction of the solution and a second fraction of said solution. The first fraction is substantially devoid of target. The solution of said first fraction contains replicatable nucleic acid templates having lower affinity than replicatable nucleic acid templates of the second fraction. The second fraction has the target or was closely associated with the target. This fraction contains replicatable nucleic acid templates having higher affinity than replicatable nucleic acid templates in the first fraction.

Turning now to Figure 4b, a second vessel 47 receives a replicatable nucleic acid template, target, reagents and polymerase. The replicatable nucleic acid template is selected for high or low affinity. Upon imposing reaction conditions, at least one subsequent generation of template is formed. The double arrow represents imposing fraction forming conditions on the target. These conditions may be imposed by chemical

or physical means, as discussed with respect to Figure 4a. The imposition of fraction forming conditions forms a first fraction of the solution and a second fraction of the solution. The first fraction is substantially devoid of target. The solution of the first fraction contains replicatable nucleic acid templates having lower affinity than replicatable nucleic acid templates of the second fraction. The second fraction has the target or was closely associated with the target. This fraction contains replicatable nucleic acid templates having higher affinity than replicatable nucleic acid templates in the first. Preferably, the formation of fractions is repeated or is performed in a continuous manner during the formation of multiple generations. Preferably, the movement of selected fractions, the imposition of fraction forming conditions, the addition of target, reagents, and polymerase is automated. Thus, replicatable nucleic acid templates exhibiting high or low affinity, or any affinity in between, can be created.

Preferably, the selection is applied in a plurality of vessels and/or in a plurality of media to reduce selection based on the composition of the vessel.

Turning now to Figure 5, a further embodiment of a device, generally designated by the numeral 51, embodying features of the present invention, is illustrated in partial cut-away. Device 51 comprises the following major elements: a housing 53, a matrix 55, and electrodes 57a-d.

Housing 53 defines a containment vessel for holding matrix 55 and a solution containing replicatable nucleic acid templates and reagents. A permeable divider 61 divides the housing into a first compartment 63 and a second compartment 65. Matrix 55 is held in a first compartment 63 or comprises one or more the walls of first compartment 63. Such walls may be pitted to form micro volumes. Second compartment 65 provides a fluid passage for fluids exiting matrix 55.

Housing 53 has an inlet 69 for receiving a solution containing one or more nucleic acid templates and other reagents and directing such solution to matrix 55. Inlet 69 is, preferably, in fluid communication with a continuous source (not shown) of templates and reagents and controlled by suitable valves and automated controls (not shown). Housing 53 has an outlet 71 for discharging fluid.

Matrix 55 comprises a packing or support material to which target or an evolvase (not shown) is immobilized. Matrix 55 allows solution to flow carrying replicatable nucleic acid templates. Under condition for replication, the templates evolve with templates with the greatest affinity for the target being retained in the matrix 55.

Electrodes 57a-d provide electromotive force to compel templates exhibiting weak affinity for the target to exit the matrix 55. Thus, electrodes 57b and c exert a weak electromotive force on templates entering the matrix 55 and electrodes 57 a and d exert a more powerful electromotive force on templates exiting the matrix 55 toward the outlet 71.

In operation, replicatable nucleic acid templates placed in the matrix 55 at the inlet 69 with a polymerase and suitable reagents. Templates are created with varying degrees of affinity to the target. Templates exhibiting weak affinity to the target are withdrawn from the matrix 55 by electromotive force created by electrodes 57a-d and conveyed from the housing 51 via second compartment 65 and outlet 71.

Templates with higher affinity to the target are retained in the matrix 55 and undergo further replication and evolution. Provided with reagents, the templates exiting outlet 71 will become constant or begin to taper off as templates evolve with such high affinity that such templates are bound to target and are unavailable to replicate. Such templates can be isolated from the target by denaturing the matrix bound target.

Figure 6 depicts a kit, generally designated by the numeral 79, for making a replicatable nucleic acid template having a selected affinity to a target. The kit comprises a housing 15 and support 17 for applying a selection to a first generation comprising at least one replicatable nucleic acid template as the replicatable nucleic acid template is replicated by a polymerase to form at least one subsequent generation comprising a replicatable nucleic acid template. The selection is based on the affinity of the replicatable nucleic acid template of different generations to the target. The polymerase introduces genetic variability between generations of the replicatable nucleic acid template to produce replicatable nucleic acid templates having different affinities to the target. The housing 15 and support also facilitate separating the replicatable nucleic acid templates on the basis of the affinity of the replicatable nucleic acid template to the target. Preferably, the support 17 has target immobilized on it surface, or is functionalized to allow an individual skilled in the art to place the target of his/ her choice on the support. The kit further comprises suitable reagents and polymerases held in appropriate vessels 81-83 (additional vessels may be required). The kit, preferably, is held in suitable packaging, such as a box 85, and provided with instructions for use.

A preferred kit comprises an evolvas.



While the preferred embodiments of the present invention have been illustrated and described, it is understood that the present invention is capable of variation and modification and therefore, should not be limited to the precise details set forth, but should  
5 include such changes and alterations that fall within the purview of the following claims.

## Claims

1. A method of making a replicatable RNA template having a selected affinity to a target,  
5 comprising the steps of:  
a.) applying a selection to a first generation comprising at least one replicatable RNA  
template as said replicatable RNA template is replicated by a RNA polymerase to form at  
least one subsequent generation comprising a replicatable RNA template, said selection  
based on the affinity of said replicatable RNA template of different generations to said  
10 target, said RNA polymerase introducing genetic variability between generations of said  
replicatable RNA template to produce replicatable RNA templates having different  
affinities to said target; and  
b.) separating the replicatable RNA templates on the basis of the affinity of said  
replicatable RNA template to the target.
- 15 2. The method of claim 1 wherein said selection is applied by passing a solution  
comprising said first and subsequent generations of said replicatable RNA template  
through or over a fixed medium, said fixed medium having said target immobilized, such  
that replicatable RNA templates having low affinity for the target and replicatable RNA  
20 templates having high affinity for the target assume different positions in said solution as  
said solution passes through said fixed medium.
3. The method of claim 2 wherein said fixed medium is provided with reagents and  
an RNA polymerase.
- 25 4. The method of claim 3 wherein said RNA polymerase is selected from the group of  
polymerases consisting of Q-Beta replicase, MS2 replicase, GA replicase, and SP  
replicase and modifications thereof.
- 30 5. The method of claim 4 wherein said RNA polymerase is non-selective with respect  
to template.
6. The method of claim 1 wherein said selection is applied by placing said first  
generation and target in a solution, and after at least one subsequent generation has been

formed, forming a first fraction of said solution and a second fraction of said solution, said first fraction substantially devoid of target, said solution of said first fraction containing replicatable RNA templates having lower affinity than replicatable RNA templates in said second fraction, said second fraction having said target containing replicatable RNA templates having higher affinity than replicatable RNA templates in said first fraction.

7. The method of claim 6 wherein said separation is performed by one or more of the processes selected from the group consisting of centrifugation, filtration, electric field electrophoresis, and precipitation.

8. The method of claim 6 wherein at least one fraction is provided with additional reagents and RNA polymerase in the presence of target.

9. The method of claim 6 wherein said selection is applied in a plurality of vessels containing said solution to reduce selection based on the composition of the vessel.

10. The method of claim 1 wherein said selection is applied by an enzyme having a catalytic entity and a target entity.

11. The method of claim 10 wherein said catalytic entity is derived from one of the RNA dependent RNA polymerases selected from the group consisting of Q-Beta replicase, MS2 replicase, GA replicase, and SP replicase and modifications thereof.

12. The method of claim 10 wherein said target is incorporated into one or more subunits of the polymerase.

13. The method of claim 10 wherein said target is covalently bonded to said catalytic entity.

14. A device for making a replicatable RNA template having a selected affinity to a target, comprising:

a.) means for applying a selection to a first generation comprising at least one replicatable RNA template as said replicatable RNA template is replicated by a RNA polymerase to

form at least one subsequent generation comprising a replicatable RNA template, said selection based on the affinity of said replicatable RNA template of different generations to said target, said RNA polymerase introducing genetic variability between generations of said replicatable RNA template to produce replicatable RNA templates having different  
5 affinities to said target; and

b.) means for separating the replicatable RNA templates on the basis of the affinity of said replicatable RNA template to the target.

10 15. The device of claim 14 wherein said means for applying a selection is a fixed medium said selection applied by passing a solution comprising said first and subsequent generations of said replicatable RNA template through said fixed medium, said fixed medium having said target immobilized, such that replicatable RNA templates having low affinity for the target and replicatable RNA templates having high affinity for the target  
15 assume different positions in said solution as said solution passes through said fixed medium.

16. The device of claim 11 wherein said fixed medium is held in a column.

20 17. A kit for making a replicatable RNA template having a selected affinity to a target, comprising:  
a.) means for applying a selection to a first generation comprising at least one replicatable RNA template as said replicatable RNA template is replicated by a RNA polymerase to form at least one subsequent generation comprising a replicatable RNA template, said  
25 selection based on the affinity of said replicatable RNA template of different generations to said target, said RNA polymerase introducing genetic variability between generations of said replicatable RNA template to produce replicatable RNA templates having different affinities to said target; and  
b.) means for separating the replicatable RNA templates on the basis of the affinity of said  
30 replicatable RNA template to the target.

18. The Kit of claim 17 wherein said RNA polymerase is evolvase.

## SEQUENCE LISTING

<110> Grossman, Abraham

InVitro Diagnostics, Inc.

<120> Methods, Compositions and Apparatus for Making Nucleic  
Acid Molecules Having a Selected Affinity to a Target  
Molecule

<130> Grossman

<140>

<141>

<150> 60/091,578

<151> 1996-07-02

<150> 1

<170> PatentIn Ver. 3.0

<210> 1

<211> 237

<212> RNA

<213> Q-beta bacteriophage

<400> 1

sdnggggacc cccccggaag ggggggacga ggugcgggca ccucguacgg gaguucgacc 60

gugacgcucu agagauuag agcacgggcu agcgcuuucg cgcucucca ggugacgcucu 120  
cgugaagagg cgcgacuccu ggcgcuuucg gugacgcacg agaaccgcca cgcugcuucg 130  
cagcguggcu ccuucggcca ggcgcugcg agaggugacc ccggaagggg gguuucc 237




FIG. 1




FIG. 3






FIG. 2








FIG. 5




FIG. 6

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/13030

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12Q 1/68; C12P 19/34

US CL : 435/6, 91.1, 91.3

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.1, 91.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, STN, caplus, medline, biosis, embase

search terms: Q beta replicase, ms2 replicase, ga replicase, sp replicase, replicatable mra template, mra polymerase

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,763,171 A (STEFANO et al) 09 June 1998, see entire document.	1-18
Y	US 5,759,773 A (TYAGI et al) 02 June 1998, see entire document.	1-18
Y	US 5,364,760 A (CHU et al) 15 November 1994, see entire document.	1-18
Y	US 5,270,163 A (GOLD et al) 14 December 1993, see entire document.	1-18
Y	US 5,472,840 A (STEFANO) 05 December 1995, see entire document.	1-18
Y	US 5,464,744 A (FARRELL et al) 07 November 1995, see entire document.	1-18

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document published on or after the international filing date

\*L\* document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\*

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*A\*

document member of the same patent family

Date of the actual completion of the international search

15 AUGUST 1999

Date of mailing of the international search report

10 SEP 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JEHANNE SOUAYA

Telephone No. (703) 308-1235

JOYCE BRIDGERS  
PARALEGAL SPECIALIST  
CHEMICAL MATRYX